# nature portfolio

| Corresponding author(s):   | Fan-Yan Wei |  |
|----------------------------|-------------|--|
| Last updated by author(s): | 2023/4/13   |  |

## **Reporting Summary**

Nature Portfolio wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Portfolio policies, see our <u>Editorial Policies</u> and the <u>Editorial Policy Checklist</u>.

| 2 | ta | t١ | S | tı | CS |
|---|----|----|---|----|----|
|   |    |    |   |    |    |

| For         | all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.  |
|-------------|--|
| n/a         | Confirmed  |
|             | $\square$ The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement  |
|             | A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly  |
|             | The statistical test(s) used AND whether they are one- or two-sided  Only common tests should be described solely by name; describe more complex techniques in the Methods section.  |
| $\boxtimes$ | A description of all covariates tested   |
|             | A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons  |
|             | A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals) |
|             | For null hypothesis testing, the test statistic (e.g. <i>F</i> , <i>t</i> , <i>r</i> ) with confidence intervals, effect sizes, degrees of freedom and <i>P</i> value noted <i>Give P values as exact values whenever suitable.</i>                        |
| $\boxtimes$ | For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings   |
| $\boxtimes$ | For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes   |
| $\boxtimes$ | Estimates of effect sizes (e.g. Cohen's <i>d</i> , Pearson's <i>r</i> ), indicating how they were calculated   |
|             | Our web collection on <u>statistics for biologists</u> contains articles on many of the points above.  |
|             |  |

#### Software and code

Policy information about availability of computer code

Data collection

No software was used for data collection.

Data analysis

GraphPad Prism (Version 9.4.1) and Microsoft excel (Version 16.66.1) were used to calculate mean, standard error and P value. Fiji software was used to analyze imaging data.

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Portfolio <u>guidelines for submitting code & software</u> for further information.

#### Data

Policy information about <u>availability of data</u>

All manuscripts must include a data availability statement. This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A description of any restrictions on data availability
- For clinical datasets or third party data, please ensure that the statement adheres to our policy

All data are available in the main text or the supplementary materials.

| Research inv       | volving hu      | man participants, their data, or biological material |
|--------------------|-----------------|--|
| Policy information | about studies v | vith   |

## **Antibodies**

Dual us

Antibodies used

Clinical data

Dual use research of concern

nti-β-Actin mAb, Code No.M177-3, MBL Phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204) Rabbit mAb, #9101, Cell Signaling p44/42 MAPK (Erk1/2) (137F5) Rabbit mAb #4695, Cell Signaling Phospho-Akt (Ser473) (D9E) XP® Rabbit mAb #4060, Cell Signaling Akt (pan) (C67E7) Rabbit mAb #4691, Cell Signaling Anti-MTCO1 antibody [1D6E1A8], ab14705, abcam TFAM (D5C8) Rabbit mAb #8076, Cell Signaling Total OXPHOS Human WB Antibody Cocktail, ab110411, abcam

Anti-FLAG antibody (M2), #F3165, Sigma

Validation

All antibodies are commercially available and were commercially validated.

## Eukaryotic cell lines

Policy information about <u>cell lines and Sex and Gender in Research</u>

For the GPCR assay, we used HEK293 cells obtained from ATCC. Cell line source(s)

Authentication The cell line used was not authenticated.

Mycoplasma contamination The cell lines was tested negative for mycoplasma contamination.

Commonly misidentified lines (See ICLAC register)

No commonly misidentified cell lines were used.

### Animals and other research organisms

Policy information about studies involving animals; ARRIVE guidelines recommended for reporting animal research, and Sex and Gender in Research

Sprague-Dawley rat pups (male and female) on postnatal day 1-3 were purchased from Japan SLC, Inc. Laboratory animals C57BL/6J mice for the isolation of adult mouse cardiomyocytes were purchased from Japan SLC Inc.

Wild animals No wild animals were used.

not applicable Reporting on sex

Field-collected samples No field-collected samples were used.

Animal studies were reviewed and approved by the ethics committees at the Tohoku University and Kyushu University and carried Ethics oversight

out in accordance with their guidelines.

Note that full information on the approval of the study protocol must also be provided in the manuscript.